

Recombinant Mouse IL-23 Receptor/IL-23R Protein (Fc Tag)

Catalog Number: PKSM041055

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Interleukin-23 receptor; IL23r; IL-23 receptor; IL-23R; interleukin 23 receptor
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Gly24-Asp372
Accession	Q8K4B4
Calculated Molecular Weight	67.4 kDa
Observed molecular weight	90-110 kDa
Tag	C-Fc

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Interleukin 23 receptor (IL23R), a heterodimer of the IL12 receptor $\beta 1$ (IL12R $\beta 1$) and IL12R $\beta 2$, is a type I cytokine receptor for IL23. IL23R is comprised of two subunits, the IL12R $\beta 1$ subunit, which is shared with several cytokines, and a subunit that is unique to IL-23. IL23, after binding to IL23R, activates memory T cells and mediates pro-inflammatory activities in part by the production of IL17 through activation of TH17 lymphocytes. IL23R is expressed on T cells, NK cells, dendritic cells, and macrophages. In fact, polymorphisms of the IL23R gene were reported to be associated with susceptibility to inflammatory diseases and autoimmune diseases such as psoriasis, multiple sclerosis, Graves's ophthalmopathy and inflammatory bowel diseases. The IL23R is known to be critically involved in the carcinogenesis of different malignant tumor.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017