Recombinant Mouse IDO1/IDO Protein (His Tag)

Catalog Number: PKSM041057



Note: Centrifuge before opening to ensure complete recovery of vial contents.

SpeciesMouseExpression HostE.coliSequenceMet1-Pro407AccessionP28776Calculated Molecular Weight47.1 kDaObserved molecular weight40-50 kDaTagN-HisPropertiesPurity> 95 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per µg of the protein as determined by the LAL method.		
Expression HostE.coliSequenceMet1-Pro407AccessionP28776Calculated Molecular Weight47.1 kDaObserved molecular weight40-50 kDaTagN-HisPropertiesPurity>95 % as determined by reducing SDS-PAGE.Endotoxin<1.0 EU per µg of the protein as determined by the LAL method.	Indole 2;3-dioxygenase;Indoleamine 2;3-dioxygenase 1;IDO-1;IDO1;IDO;INDO	
SequenceMet1-Pro407AccessionP28776Calculated Molecular Weight47.1 kDaObserved molecular weight40-50 kDaTagN-HisPropertiesPurity> 95 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per µg of the protein as determined by the LAL method.	Mouse	
AccessionP28776Calculated Molecular Weight47.1 kDaObserved molecular weight40-50 kDaTagN-HisPropertiesPurity> 95 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per μg of the protein as determined by the LAL method.StorageStore at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.ShippingThis product is provided as liquid. It is shipped at frozen temperature with b	E.coli	
Calculated Molecular Weight47.1 kDaObserved molecular weight40-50 kDaTagN-HisPropertiesPurity>95 % as determined by reducing SDS-PAGE.Endotoxin<1.0 EU per μg of the protein as determined by the LAL method.	Met1-Pro407	
Observed molecular weight40-50 kDaTagN-HisPropertiesPurity> 95 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per μg of the protein as determined by the LAL method.	P28776	
TagN-HisPropertiesPurity> 95 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per μg of the protein as determined by the LAL method.	47.1 kDa	
Properties Purity > 95 % as determined by reducing SDS-PAGE. Endotoxin < 1.0 EU per μg of the protein as determined by the LAL method. Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles. Shipping This product is provided as liquid. It is shipped at frozen temperature with b	40-50 kDa	
Purity> 95 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per μg of the protein as determined by the LAL method.		
Endotoxin< 1.0 EU per μg of the protein as determined by the LAL method.		
StorageStore at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.	> 95 % as determined by reducing SDS-PAGE.	
Shipping This product is provided as liquid. It is shipped at frozen temperature with b	< 1.0 EU per μ g of the protein as determined by the LAL method.	
	Store at $< -20^{\circ}$ C, stable for 6 months. Please minimize freeze-thaw cycles.	
ice/gel packs. Upon receipt, store it immediately at $< -20^{\circ}$ C.	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20° C.	
FormulationSupplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 500mM NaCl, 0.0Tween 80, 1mM EDTA, 50% Glycerol,pH 8.0.	Supplied as a 0.2 μ m filtered solution of 20mM Tris-HCl, 500mM NaCl, 0.01% Tween 80, 1mM EDTA, 50% Glycerol,pH 8.0.	
Reconstitution Not Applicable	Not Applicable	
Data		

kDa	MK	R
120 90		
60		
40		-
30	-	ana angle
20	-	
14	-	

> 95 % as determined by reducing SDS-PAGE.

Background

Indoleamine 2,3-dioxygenase (IDO) is a heme enzyme that initiates the oxidative degradation of the least abundant, essential amino acid, l-tryptophan, along the kynurenine pathway. This protein is normally expressed in the dendritic cells, macrophages, microglia, eosinophils, fibroblasts, endothelial cells, and most tumor cells. IDO activity is associated with immunosuppression and immune attenuation. Several studies showed that IDO can contribute to immune escape when expressed directly in tumor cells or when expressed in immunosuppressive antigen presenting cells such as tolerogenic dendritic cells or tumor associated macrophages. IDO also is a promising therapeutic target for the treatment of cancer, chronic viral infections, and other diseases characterized by pathological immune suppression.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com