Recombinant Mouse Interleukin-10/IL-10 Protein

Catalog No. PKSM041066

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Interleukin-10;II10;IL-10;Cytokine synthesis inhibitory factor;CSIF;
Species	Mouse
Expression Host	E.coli
Sequence	Ser19-Ser178
Accession	P18893
Calculated Molecular Weight	19.7 kDa
Observed molecular weight	19 kDa
Tag	C-His
Properties	
Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.1 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	

kDa 75-63-48-35-25-17-11-

> 98 % as determined by reducing SDS-PAGE.

Background

Mouse II10 is the prototypic member of the IL-10 cytokine family, including IL-10, IL-19, IL-20, IL-22 (IL-TIF), IL-24 and IL-26. Many viruses encode viral members of the IL-10 family, such as Epstein-Barr virus (EBV) and human cytomegalovirus (HCMV). Its main function is inhibiting the synthesis of a number of cytokines, including IFN-gamma,

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

IL-2, IL-3, TNF and GM-CSF produced by activated macrophages and by helper T-cells. Although human and mouse IL-10 are 81% identical at the nucleotide and amino acid level, mouse IL-10 is species-specific and does not act on human cells. Interestingly, Human IL-10 is active on mouse cells.

For Research Use Only