

Recombinant Mouse Interleukin-13/IL-13 Protein (His Tag)

Catalog No. PKSM041074

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Interleukin-13;IL-13;T-Cell Activation Protein P600;IL13;IL-13
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Ser26-Phe131
Accession	P20109
Calculated Molecular Weight	12.7 kDa
Observed molecular weight	15-30 kDa
Tag	C-His
Bioactivity	Measured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED ₅₀ for this effect is 2-13 ng/ml.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Mouse interleukin 13 (mIL-13) is a pleiotropic cytokine produced by activated Th2 cells. IL-13 induces B cell

For Research Use Only

proliferation and immunoglobulin production. It contains a four helical bundle with two internal disulfide bonds. Mouse IL13 shares 58% sequence identity with human protein and exhibits cross-species activity. IL13 signals via receptor IL13R (type2, IL4R) and activates STAT-6. IL13 initially binds IL-13R α 1 with low affinity and triggers association of IL4R α , generating a high affinity heterodimeric receptor IL13R and eliciting downstream signals. IL13 also binds IL-13R α 2 with high affinity, which plays a role in a negative feedback system of IL13 signaling. IL13 is an important mediator of allergic inflammation and disease.