

Recombinant Mouse Interleukin-33/IL-33 Protein

Catalog No. PKSM041090

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Interleukin 33;IL-33;IL33;C9orf26;NKHEV;Interleukin-1 family member 11;DVS27;NF-HEV and IL- 1F11
Species	Mouse
Expression Host	E.coli
Sequence	Ser109-Ile266
Accession	Q8BVZ5
Calculated Molecular Weight	18.5 kDa
Observed molecular weight	17-25 kDa
Tag	C-His

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.1 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

Mouse Interleukin 33 (IL-33) is a 30 kDa proinflammatory cytokine which may also regulates gene transcription in producer cells. IL-33 is constitutively expressed in smooth muscle and airway epithelia. IL-33 was identified based on

For Research Use Only

sequence and structural homology with IL-1 family cytokines. It is up-regulated in arterial smooth muscle, dermal fibroblasts, and keratinocytes following IL-1 alpha or IL-1 beta stimulation. IL-33 is structurally related to IL-1, which induces helper T cells to produce type 2 cytokines and acts through the receptor IL1RL-1. Binding IL-33 to this receptor activates NF-kappa-B and MAP kinases and induces in vitro Th2 cells to produce cytokines. In vivo, IL-33 induces the expression of IL-4, IL-5, IL-13 and also leads to severe pathological changes in mucosal organs.