Recombinant Mouse LAG3 Protein (His Tag)

Catalog No. PKSM041104

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	Lymphocyte activation gene 3 protein;CD223;Lag3	
Species	Mouse	
Expression Host	HEK293 Cells	
Sequence	Ser23-Leu442	
Accession	Q61790	
Calculated Molecular Weight	46.2 kDa	
Observed molecular weight	55-80 kDa	
Tag	C-His	
Bioactivity	Not validated for activity	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	

Data

kDa	MK	R
120 90	-	
60	- 1	-
40	-	11.74
30	-	
20	•	
14	12	

> 95 % as determined by reducing SDS-PAGE.

Background

Lymphocyte-activation gene 3 (LAG3), also known as CD223, is a type I transmembrane protein with four extracellular Ig-like domains, designated D1 to D4 and belongs to the immunoglobulin superfamily. The gene for LAG3 lies adjacent

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

to the gene for CD4 on human chromosome 12p13.32 and shares approximately 20% identical to the CD4 gene. LAG3 is expressed on activated T cells, natural killer cells, B cells and plasmacytoid dendritic cells. LAG3 binds with high affinity to MHC class II molecules, and it interferes competitively with the binding of CD4 to MHC class II and thereby blocks the transduction of stimulatory signals mediated by this interaction. LAG3 negatively regulates cellular proliferation, activation, and homeostasis of T cells, and plays an important role in Treg suppressive function. LAG3 is the target of various drug development programs to develop new treatments for cancer and autoimmune disorders. The soluble form, sLAG-3, is being developed as a cancer drug.

For Research Use Only