

Recombinant Mouse MBL2/MBP-C Protein

Catalog No. PKSM041108

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Mannose binding lectin (C);isoform CRA_b;Mannose-binding protein C;Mbl2;MBL-2;Mannose Binding Lectin 2
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Glu19-Asp244
Accession	Q3UEK1
Calculated Molecular Weight	24.0 kDa
Observed molecular weight	28 kDa
Tag	None
Bioactivity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Mannose-binding Lectin (MBL) is an acute phase protein bearing to the family of collectins produced by the liver as a

For Research Use Only

monomer that forms a triple helix. Once released in serum, it further polymerizes forming dimers to octamers. The degree of serum polymerization is critical for the biological activity of MBL. MBL has higher affinity to microbial polysaccharides or their glycoconjugates. MBL was shown earlier to bind cell surfaces of bacteria, fungi, protozoa and viruses and acts as an acute-phase plasma protein (APP) during infection and inflammation. MBL activates the lectin-complement pathway, promotes opsonophagocytosis and modulates inflammation.