## Recombinant Mouse TNFSF4/OX40L Protein (His Tag)

Catalog No. PKSM041119

*Note:* Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Synonyms	Tumor necrosis factor ligand superfamily member 4;OX40 ligand;OX40L;CD252;Tnfsf4		
Species	Mouse		
Expression Host	HEK293 Cells		
Sequence	Ser51-Leu198		
Accession	P43488		
Calculated Molecular Weight	17.7 kDa		
Observed molecular weight	20-23 kDa		
Tag	N-His		
Bioactivity	Not validated for activity		
Properties			
Purity	> 95 % as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU per $\mu$ g of the protein as determined by the LAL method.		
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.		
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.		
Reconstitution	Please refer to the printed manual for detailed information.		
Data			

kDa	MK	R	
120 90			
60			3
40	المنت		
30	-		
20	-"		
14	-151		

> 95 % as determined by reducing SDS-PAGE.

## Background

OX40 ligand (OX40L), also called CD252, is a single-pass type II membrane protein of the TNF/TNF receptor

## For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u>

## **Elabscience**®

superfamily. OX40L is expressed by DCs, macrophages and B cells and signals via its cognate receptor OX40 which is mainly expressed on APCs. OX40L/OX40 interactions are important in T-cell activation and survival and for the generation of memory T cells from activated effector T cells. OX40L–OX40 co-stimulation leads to activation of TNF receptor associated factor (TRAF) 2, 3 and 5. This pathway has been shown to prolong the survival of effector CD4+Th cells as well as contributes to generation of memory T cells.

For Research Use Only