

Recombinant Mouse RANK/TNFRSF11A Protein (His Tag)

DIA • AN[®]
by Elabscience

Catalog Number:PKSM041130

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Receptor activator of NF- κ B;tumor necrosis factor receptor superfamily member 11A;TRANCE receptor;Osteoclast differentiation factor receptor;NF κ B activator;TRANCER;CD265;TNFRSF11A;TRANCE R;CD265 antigen;ODFR
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Val31-Ser214
Accession	O35305
Calculated Molecular Weight	21.3 kDa
Observed molecular weight	26-30 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Receptor activator of NF- κ B(RANK,TNFRSF11A) belongs to one member of tumor necrosis factor receptor family.It is a receptor for TNFSF11/RANKL/TRANCE/OPGL. This gene encodes a type 1 membrane protein with a 30 amino acids (aa) signal peptide, 184 aa extracellular region , a 20 aa transmembrane domain and a 391 aa cytoplasmic region. Human and murine RANK share 81% aa identity in their extracellular domains. RANK is ubiquitous highly expressed in trabecular bone, thymus, small intestine, lung, brain and kidney, but weakly expressed in spleen and bone marrow. After binding its ligand RANKL, RANK can activate signaling pathways such as NF- κ B, JNK, ERK, p38, and Akt/PKB,

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Mouse RANK/TNFRSF11A Protein (His Tag)

Catalog Number:PKSM041130

DIA • AN®
by Elabscience

through TRAF protein phosphorylation. RANK/TNFRSF11A signaling is largely considered to be growth promoting and apoptosis reducing such as the effects observed in osteoclasts. RANK/TNFRSF11A was also found to be involved in the regulation of interactions between T-cells and dendritic cells.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com