

Recombinant Mouse SerpinD1/HCF2 Protein (His Tag)

Catalog No. PKSM041140

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Heparin cofactor 2;Heparin cofactor II;HC-II;Protease inhibitor leuserpin-2;Serpin D1
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Glu24-Ser478
Accession	P49182
Calculated Molecular Weight	53.1 kDa
Observed molecular weight	70-90 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 50mM Tris-HCl, 150mM NaCl, 5% Mannitol, 0.06% Tween 80, pH8.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

SerpinD1, also known as heparin cofactor II(HC-II), is a member of Serpin superfamily of the serine proteinase inhibitors. It is a single chain glycoprotein with a size of 66.5 kDa and is secreted from hepatocytes. HC-II acts as a thrombin inhibitor in the coagulation cascade, in a glycosaminoglycan-dependent pathway using the release of a sequestered hirudin-like N-terminal tail for interaction with thrombin. This serpin belongs to multiple member group V2 of vertebrate serpin classification. It has been suggested that HC-II is a predictor of decreased atherosclerosis in the elderly and protective against atherosclerosis in mice. HCII can used as a predictive biomarker and therapeutic target for atherosclerosis.