

# Recombinant Mouse RANKL/TNFSF11 Protein

Catalog Number:PKSM041166



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

|                                    |   |
|------------------------------------|---|
| <b>Synonyms</b>                    | Tumor necrosis factor ligand superfamily member 11;Tnfsf11;Osteoclast differentiation factor;ODF;Osteoprotegerin ligand;OPGL;Receptor activator of nuclear factor kappa-B ligand;RANKL;TNF-related activation-induced cytokine;TRANCE;CD254 |
| <b>Species</b>                     | Mouse   |
| <b>Expression Host</b>             | E.coli  |
| <b>Sequence</b>                    | Lys158-Asp316   |
| <b>Accession</b>                   | O35235  |
| <b>Calculated Molecular Weight</b> | 17.9 kDa  |
| <b>Observed molecular weight</b>   | 18 kDa  |
| <b>Tag</b>                         | No tag  |

## Properties

|                       |   |
|-----------------------|---|
| <b>Purity</b>         | > 95 % as determined by reducing SDS-PAGE.  |
| <b>Endotoxin</b>      | < 0.01 EU per µg as determined by the LAL method.   |
| <b>Storage</b>        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| <b>Shipping</b>       | This product is provided as lyophilized powder which is shipped with ice packs.   |
| <b>Formulation</b>    | Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.   |
| <b>Reconstitution</b> | Please refer to the printed manual for detailed information.  |

## Data



## Background

Mouse tumor necrosis factor ligand superfamily member 11 (Tnfsf11) is a member of the tumor necrosis factor (TNF) cytokine family. Tnfsf11 is widely expressed in cells including T cells and T cell rich organs, such as thymus and lymph nodes. This cytokine can bind to TNFRSF11B/OPG and TNFRSF11A/RANK. Tnfsf11 is involved in a number of fundamental biological processes such as acting as regulator of interactions between T-cells and dendritic cells, the regulation of the T-cell-dependent immune response and enhancing bone-resorption in humoral hypercalcemia of malignancy. It augments the ability of dendritic cells to stimulate naive T-cell proliferation.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017