# **Recombinant Mouse TGFBR2 Protein (His Tag)**

Catalog Number: PKSM041169



Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

**Synonyms** TGF-beta receptor type-2;TGFR-2;TGF-beta type II receptor;Transforming growth

factor-beta receptor type II;TGF-beta receptor type II;TbetaR-II;Tgfbr2

Species Mouse

Expression Host

Sequence

Ile24-Asp159

Accession

Q62312-2

Calculated Molecular Weight

Observed molecular weight

Tag

HEK293 Cells

Ile24-Asp159

Q62312-2

25-38 kDa

C-His

**Bioactivity** Measured by its ability to inhibit TGF-beta 1 activity on TF-1 human

erythroleukemic cells. The ED<sub>50</sub> for this effect is 1680.39 ng/ml in the presence of

1ng/ml of recombinant human TGF-beta 1.

## **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

# **Background**

Transforming growth factor- $\beta$  (TGF- $\beta$ ) is an essential regulator in the processes of development, cell proliferation, and extracellular matrix deposition. TGF- $\beta$  regulates cellular processes by binding to three high-affinity cell surface receptors: TGF- $\beta$  receptor type I (TGF- $\beta$ -RII), TGF- $\beta$  receptor type II (TGF- $\beta$ -RII), and TGF- $\beta$  receptor type III (TGF- $\beta$ -RIII). TGF- $\beta$  RII is consists of a C-terminal protein kinase domain and an N-terminal ectodomain and belongs to transforming

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

# **Recombinant Mouse TGFBR2 Protein (His Tag)**

Catalog Number: PKSM041169



growth factor-beta (TGF- $\beta$ ) receptor subfamily. TGF- $\beta$  RII has a protein kinase domain which can form a heterodimeric complex with another receptor protein and bind TGF-beta. This receptor/ligand complex phosphorylates protein will enter the nucleus and regulate the transcription of a subset of genes related to cell proliferation.

### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>