



A Reliable Research Partner in Life Science and Medicine

# Purified Anti-Mouse CD3 Antibody[17A2]

Catalog No. E-AB-F1013A Reactivity Mouse Storage Store at 2~8°C, Avoid freeze / thaw cycles **Applications FCM** 

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### **Antigen Information**

**Alternate Names** T-cell surface glycoprotein CD 3epsilon/delta/gamma/zeta

chain,CD3E/D/G/Z,CD3e/d/g/z,CD3E/D/G/Z,CD3

**Uniprot ID** P04235,P11942,P22646,P24161

**Background** CD3, also known as T3, is a member of the Ig superfamily and primarily expressed on T cells,

> NK-T cells, and at different levels on thymocytes during T cell differentiation. CD3 is composed of CD3 $\epsilon$ ,  $\delta$ ,  $\gamma$  and  $\zeta$  chains. It forms a TCR complex by associating with TCR  $\alpha/\beta$  or  $\gamma/\delta$  chains. CD3 plays a critical role in TCR signal transduction, T cell activation, and antigen recognition by

binding the peptide/MHC antigen complex.

#### **Product Details**

Form Liquid Concentration 0.5 mg/mL Size  $25 \mu g / 100 \mu g$ Clone No. 17A2 Host Rat

Rat IgG2b, κ **Isotype** Reactivity Mouse **Application FCM** 

**Isotype Control** Purified Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09843A]

**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Biological ice pack at 4 °C Shipping Stability & Storage Keep as concentrated solution. Store at 2~8°C .Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Toll-free: 1-888-852-8623 Email: techsupport@elabscience.com Web: www.elabscience.com

Tel: 1-832-243-6086

Fax: 1-832-243-6017

#### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

## **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 0.25~\mu g$  per  $10^6$  cells in  $100~\mu L$  volume or  $100~\mu L$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

### **Related Information**

- 1. Sample Preparation for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5594.html">https://www.elabscience.com/List-detail-5594.html</a>
- 2. Staining Cell Surface Targets for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5568.html">https://www.elabscience.com/List-detail-5568.html</a>
- 3. Flow Cytometry Troubleshooting Tips <a href="https://www.elabscience.com/List-detail-5593.html">https://www.elabscience.com/List-detail-5593.html</a>
- 4. How to select the appropriate detection channel through the spectrogram? <a href="https://www.elabscience.com/List-detail-459742.html">https://www.elabscience.com/List-detail-459742.html</a>

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com