Elabscience®

Biotin Anti-Mouse CD49b/pan-NK cells Antibody[DX5]

Catalog No.E-AB-F1116BStorageStore at 2~8°C, Avoid freeze / thaw cycles

ReactivityMouseApplicationsFCM

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

| Alternate Names | Integrin alpha-2,CD49 antigen-like family member B,Collagen receptor,Platelet membrane glycoprotein Ia,GPIa,VLA-2 subunit alpha,CD49b |
|-----------------|---|
| Uniprot ID | Q62469 |
| Background | DX5 antigen has been recently characterized as CD49b. It is a 150 kD integrin α chain also |
| | known as $\alpha 2$ integrin, VLA-2 α chain, and integrin $\alpha 2$ chain. CD49b non-covalently associates |
| | with CD29 (\beta1 integrin) to form the CD49b/CD29 complex known as VLA-2, a receptor for |
| | collagen and laminin. CD49b is expressed on platelets, the majority of NK cells, NKT cells, and a |
| | small subset of CD8+ T cells (this population can be significantly increased following viral |
| | infection). DX5 is used for the identification and isolation of NK cells, and is especially useful |
| | for identifying NK cells in mice lacking the NK1.1 antigen. |

Product Details

| Form | Liquid |
|---------------------|---|
| Concentration | 0.5 mg/mL |
| Size | 25µg/100µg |
| Clone No. | DX5 |
| Host | Rat |
| Isotype | Rat IgM, ĸ |
| Reactivity | Mouse |
| Application | FCM |
| Isotype Control | Biotin Rat IgM, κ Isotype Control[RTK2118] [Product E-AB-F09773B] |
| Storage Buffer | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant. |
| Shipping | Biological ice pack at 4 °C |
| Stability & Storage | Keep as concentrated solution. |
| | Store at 2~8°C .Do not freeze. |
| | This product is guaranteed up to one year from purchase. |

For Research Use Only

Elabscience®

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \ \mu g$ per 10⁶ cells in 100 μL volume or 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? <u>https://www.elabscience.com/List-detail-459742.html</u>

For Research Use Only