



A Reliable Research Partner in Life Science and Medicine

# Biotin Anti-Mouse CD132 Antibody[3E12]

Catalog No. E-AB-F1015B Reactivity Mouse Storage Store at 2~8°C, Avoid freeze / thaw cycles **Applications FCM** 

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### **Antigen Information**

**Alternate Names** Cytokine receptor common subunit gamma,II2rg,Interleukin-2 receptor subunit gamma,IL-2

receptor subunit gamma, IL-2R subunit gamma, IL-2RG, gammaC, p64, CD132

**Uniprot ID** P34902

**Background** CD132 is a 64-70 kD type I transmembrane glycoprotein of the Ig superfamily, and is also known

> as common γ chain (γc) or IL-2 receptor γ subunit. It is expressed broadly on T- and Blymphocytes, NK cells, monocytes, and granulocytes. CD132 is an essential component of cytokine receptors for IL-2, IL-4, IL-7, IL-9, IL-15, and IL-21. Ligand binding induces tyrosine phosphorylation and initiates signaling through a JAK/STAT pathway. CD132 mutation results in

X-linked severe combined immune deficiency (XSCID).

#### **Product Details**

Form Liquid Concentration 0.5 mg/mL $25 \mu g / 100 \mu g$ Size Clone No. 3E12 Host Rat

Rat IgG2b, κ **Isotype** Reactivity Mouse **Application FCM** 

**Isotype Control** Biotin Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09843B]

Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant. **Storage Buffer** 

**Shipping** Biological ice pack at 4 °C Stability & Storage Keep as concentrated solution.

Store at 2~8°C. Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Email: techsupport@elabscience.com Web: www.elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086

Fax: 1-832-243-6017

### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

## **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 1.0 \,\mu g$  per  $10^6$  cells in  $100 \,\mu L$  volume or  $100 \,\mu L$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

### **Related Information**

- 1. Sample Preparation for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5594.html">https://www.elabscience.com/List-detail-5594.html</a>
- 2. Staining Cell Surface Targets for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5568.html">https://www.elabscience.com/List-detail-5568.html</a>
- 3. Flow Cytometry Troubleshooting Tips <a href="https://www.elabscience.com/List-detail-5593.html">https://www.elabscience.com/List-detail-5593.html</a>
- 4. How to select the appropriate detection channel through the spectrogram? <a href="https://www.elabscience.com/List-detail-459742.html">https://www.elabscience.com/List-detail-459742.html</a>

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017