



A Reliable Research Partner in Life Science and Medicine

# Biotin Anti-Mouse CD40 Antibody[FGK4.5/FGK45]

Catalog No. E-AB-F1028B Reactivity Mouse Storage Store at 2~8°C, Avoid freeze / thaw cycles **Applications FCM** 

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Antigen Information**

**Alternate Names** Tumor necrosis factor receptor superfamily member 5,Cd40,B-cell surface antigen

CD40,Bp50,CD40L receptor,CD40,Tnfrsf5

**Uniprot ID** P27512

**Background** CD40 is a 48 kD type I transmembrane glycoprotein also known as Bp50. It is a member of the

tumor necrosis factor receptor (TNFR) superfamily and is expressed on B cells, basal epithelial

cells, macrophages, follicular dendritic cells, endothelial cells, and a subset of CD34+

hematopoietic progenitors. CD40 regulates B cell development/maturation, Ig isotype switching and, in combination with other signals such as IL-4, protects B cells from surface Ig-induced apoptosis and promotes proliferation. Interaction of CD40 with its ligand CD154 (gp39), which is

expressed on activated T cells, is important in costimulation and immune regulation.

### **Product Details**

**Form** Liquid Concentration 0.5 mg/mL Size 25µg/100µg Clone No. FGK4.5/FGK45

Host Rat

Isotype Rat IgG2a, κ Reactivity Mouse **FCM Application** 

**Isotype Control** Biotin Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833B]

**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping Biological ice pack at 4 °C Stability & Storage Keep as concentrated solution.

Store at 2~8°C .Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Toll-free: 1-888-852-8623 Email: techsupport@elabscience.com Web: www.elabscience.com

Tel: 1-832-243-6086 Fax: 1-832-243-6017

#### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

## **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 1.0 \,\mu g$  per  $10^6$  cells in  $100 \,\mu L$  volume or  $100 \,\mu L$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

### **Related Information**

- 1. Sample Preparation for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5594.html">https://www.elabscience.com/List-detail-5594.html</a>
- 2. Staining Cell Surface Targets for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5568.html">https://www.elabscience.com/List-detail-5568.html</a>
- 3. Flow Cytometry Troubleshooting Tips <a href="https://www.elabscience.com/List-detail-5593.html">https://www.elabscience.com/List-detail-5593.html</a>
- 4. How to select the appropriate detection channel through the spectrogram? <a href="https://www.elabscience.com/List-detail-459742.html">https://www.elabscience.com/List-detail-459742.html</a>

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017