



A Reliable Research Partner in Life Science and Medicine

Biotin Anti-Mouse CD40 Antibody[FGK4.5/FGK45]

Catalog No. E-AB-F1028B Reactivity Mouse Storage Store at 2~8°C, Avoid freeze / thaw cycles **Applications FCM**

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names Tumor necrosis factor receptor superfamily member 5,Cd40,B-cell surface antigen

CD40,Bp50,CD40L receptor,CD40,Tnfrsf5

Uniprot ID P27512

Background CD40 is a 48 kD type I transmembrane glycoprotein also known as Bp50. It is a member of the

tumor necrosis factor receptor (TNFR) superfamily and is expressed on B cells, basal epithelial

cells, macrophages, follicular dendritic cells, endothelial cells, and a subset of CD34+

hematopoietic progenitors. CD40 regulates B cell development/maturation, Ig isotype switching and, in combination with other signals such as IL-4, protects B cells from surface Ig-induced apoptosis and promotes proliferation. Interaction of CD40 with its ligand CD154 (gp39), which is

expressed on activated T cells, is important in costimulation and immune regulation.

Product Details

Form Liquid Concentration 0.5 mg/mL Size 25µg/100µg Clone No. FGK4.5/FGK45

Host Rat

Isotype Rat IgG2a, κ Reactivity Mouse **FCM Application**

Isotype Control Biotin Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833B]

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping Biological ice pack at 4 °C Stability & Storage Keep as concentrated solution.

Store at 2~8°C .Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Toll-free: 1-888-852-8623 Email: techsupport@elabscience.com Web: www.elabscience.com

Tel: 1-832-243-6086 Fax: 1-832-243-6017

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \,\mu g$ per 10^6 cells in $100 \,\mu L$ volume or $100 \,\mu L$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/List-detail-459742.html

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017