

Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

Biotin Anti-Mouse CD120b/TNFR2 Antibody[TR75-54.7]

Catalog No. E-AB-F1035B Reactivity Mouse Storage Store at 2~8°C, Avoid freeze / thaw cycles Applications FCM

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names Tumor necrosis factor receptor superfamily member 1B, Tnfrsf1b, Tumor necrosis factor receptor

2,TNF-R2,TNF-RII,TNFR-II,p75,p80 TNF-alpha receptor,CD120b

Uniprot ID P25119

Background CD120b is a 75 kD type I transmembrane protein, also known as Tumor Necrosis Factor

Receptor Type II (TNFRII) or p75. It is expressed on a variety of cells at low levels; the

expression is upregulated upon activation. This receptor binds both TNF- α and LT- α (also known as TNF- β). In association with TRAF1 and TRAF2, the receptor crosslinking induced by TNF- α or LT- α trimers is critical for signal transduction, leading to apoptosis, NF-kB activation,

increased expression of proinflammatory genes, tumor necrosis, and cell differentiation

depending on cell type and differentiation state.

Product Details

Host Armenian Hamster **Isotype** Armenian Hamster IgG

Reactivity Mouse **Application** FCM

Isotype Control Biotin Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09853B]

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping Biological ice pack at 4 °C **Stability & Storage** Keep as concentrated solution.

Store at 2~8°C .Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \,\mu g$ per 10^6 cells in $100 \,\mu L$ volume or $100 \,\mu L$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/List-detail-459742.html

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017