



A Reliable Research Partner in Life Science and Medicine

Biotin Anti-Human/Mouse CD44 Antibody[IM7]

Catalog No. E-AB-F1100B Reactivity Human, Mouse

Storage Store at 2~8°C, Avoid freeze / thaw cycles **Applications FCM**

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names CD44 antigen, CD44, CDw44, Epican, Phagocytic glycoprotein 1, PGP-1, Phagocytic glycoprotein

I,PGP-I,CD44,LHR, MDU2, MDU3, MIC4

Uniprot ID P15379,P16070

Background CD44 is a 80-95 kD glycoprotein also known as Hermes, Pgp1, H-CAM, or HUTCH. It is

> expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. As B and T cells become activated or progress to the memory stage, CD44 expression increases from low or mid levels to high levels. Thus, CD44 has been reported to be a valuable marker for memory cell subsets. High CD44 expression on Treg cells has been associated with potent suppressive function via high production of IL-10. CD44 is an adhesion molecule involved in leukocyte attachment to

and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of

inflammation, and leukocyte aggregation.

Product Details

Form Liquid Concentration 0.5 mg/mL Size $25\mu g/100\mu g$

Clone No. IM7 Host Rat

Isotype Rat IgG2b, κ Human, Mouse Reactivity

Application FCM

Biotin Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09843B] **Isotype Control**

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping Biological ice pack at 4 °C Stability & Storage Keep as concentrated solution.

Store at 2~8°C. Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \,\mu g$ per 10^6 cells in $100 \,\mu L$ volume or $100 \,\mu L$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/List-detail-459742.html

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017