



A Reliable Research Partner in Life Science and Medicine

Recombinant Rat HER2/ErbB2 Protein (aa 67-323, His Tag)

Catalog No. PKSR030455

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Receptor tyrosine-protein kinase erbB-2; Epidermal growth factor receptor-related

protein;Proto-oncogene Neu;Proto-oncogene c-ErbB-2;p185erbB2;p185neu;CD340; ERBB2;ENV;ENVW;ERVWE1;HER-2;HER-2/neu;HER2;HERV-7q;HERV-W-

ENV;HERV7Q;HERVW;HERVWENV;MLN 19;MLN19

Species Rat
Expression Host E.coli

Sequence Ala67-Val323

AccessionP06494Calculated Molecular Weight29.3 kDaObserved molecular weight32 kDaTagC-His

Bioactivity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of PBS, 5% Trehalose, 4M Urea, pH

7.4.

Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Background

ERBB2 belongs to the protein kinase superfamily, Tyr protein kinase family and EGF receptor subfamily. It contains a protein kinase domain. ERBB2 is widely expressed in epithelial cells, and amplification and/or overexpression of ErbB2 has been reported associated with malignancy and a poor prognosis in numerous carcinomas, including breast, prostate and ovarian cancers. Rat ERBB2 is an essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. ErbB2 mediates signalling pathways which involve mitogen-activated protein kinase and phosphatidylinositol-3 kinase, this receptor plays a key role in development, cell proliferation and differentiation.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com