## Recombinant Rat CD86/B7-2 Protein (His Tag)

Catalog Number: PKSR030456



Note: Centrifuge before opening to ensure complete recovery of vial contents.

**Description** 

Synonyms T-lymphocyte activation antigen CD86; Activation B7-2

antigen;CD86;B7-2;B7.2;B70;CD28LG2;LAB72

**Species** Rat

Expression Host HEK293 Cells
Sequence Vla29-Lys247

AccessionO35531Calculated Molecular Weight25.9 kDaObserved molecular weight35-55 kDaTagC-His

**Properties** 

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin**  $< 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$ 

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## **Background**

T-lymphocyte activation antigen CD86 (B7-2) is a glycosylated protein in the B7 family. B7 family members are transmembrane cell surface molecules that play important roles in immune activation and the maintenance of immune tolerance. It is highly expressed on activated antigen presenting cells. CD86 involved in the costimulatory signal essential for T-lymphocyte proliferation and interleukin-2 production, by binding CD28 or CTLA-4. It may play a critical role in the early events of T-cell activation and costimulation of naive T-cells, such as deciding between immunity and anergy that is made by T-cells within 24 hours after activation. It is expressed by activated B-lymphocytes and monocytes and promoted by MARCH8 and results in endocytosis and lysosomal degradation.

## For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com