

Recombinant Human Interleukin-12/IL-12 Protein

Catalog No. PKSH033284

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CLMF;CLMF2;IL-12A;IL-12B;IL12;IL12 p70;IMD28;IMD29;NFSK;NKSF;NKSF1;NKSF2;P35
Species	Human
Expression Host	HEK293 Cells
Sequence	Arg23-Ser219&Ile23-Ser328
Accession	P29459&P29460
Calculated Molecular Weight	22.5&34.7 kDa
Observed molecular weight	25-38&40-50 kDa
Tag	None
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

IL-12 is a heterodimeric pleiotropic cytokine made up of a 40 kDa (p40) subunit and a 35 kDa (p35) subunit. Human and

For Research Use Only

mouse IL-12 share 70% and 60% amino acid sequence identity in their p40 and p35 subunits, respectively. IL-12 is involved in the differentiation of naive T cells into Th1 cells. It is known as a T cell-stimulating factor, which can stimulate the growth and function of T cells. It stimulates the production of interferon-gamma (IFN- γ) and tumor necrosis factor-alpha (TNF- α) from T cells and natural killer (NK) cells, and reduces IL-4 mediated suppression of IFN- γ . T cells that produce IL-12 have a coreceptor, CD30, which is associated with IL-12 activity. IL-12 plays an important role in the activities of natural killer cells and T lymphocytes. IL-12 mediates enhancement of the cytotoxic activity of NK cells and CD8+ cytotoxic T lymphocytes.