A Reliable Research Partner in Life Science and Medicine

Recombinant Human MAX Protein (His Tag)

Catalog No. PKSH033317

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Protein Max;Class D Basic Helix-Loop-Helix Protein 4;bHLHd4;;Myc-Associated

Factor X;MAX;BHLHD4

SpeciesHumanExpression HostE.coli

SequenceMet 1-Ser151AccessionP61244-2Calculated Molecular Weight18.3 kDaObserved molecular weight21 kDaTagC-His

Bioactivity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per μg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 250mM NaCl, pH

8.5.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Myc-Associated Factor X (MAX) is a member of the basic helix-loop-helix leucine zipper (bHLHZ) family of transcription factors. It contains 1 basic helix-loop-helix (bHLH) domain. It is found in the brain, heart, and lung at high levels while lower levels are seen in the liver, kidney, and skeletal muscle. MAX forms a sequence-specific DNA-binding protein complex with MYC or MAD which recognizes the core sequence 5'-CAC[GA]TG-3'. The MYC-MAX complex is a transcriptional activator, whereas the MAD-MAX complex is a repressor. It may repress transcription via the recruitment of a chromatin remodeling complex containing H3 'Lys-9' histone methyltransferase activity.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com