
Recombinant Human OLFM4 Protein (His Tag)

Catalog Number: PKSH033360



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Synonyms | Olfactomedin-4;OLM4;Antiapoptotic protein GW112;G-CSF-stimulated clone 1 protein;hGC-1;hOLfD;OLFM4;GW112 |
|-----------------------------|---|
| Species | Human |
| Expression Host | HEK293 Cells |
| Sequence | Asp21-Gln510 |
| Accession | Q6UX06 |
| Calculated Molecular Weight | 56.9 kDa |
| Observed molecular weight | 60-90 kDa |
| Tag | C-His |
| Properties | |
| Purity | > 80 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per μ g of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 µm filtered solution of PBS, 5% Trehalose, PH7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |
| Data | |



> 80 % as determined by reducing SDS-PAGE.

Background

Olfactomedin-4/OLFM4 is a secreted protein which contains one olfactomedin-like domain. OLFM4 is expressed during myeloid lineage development; it strongly expressed in the prostate; small intestine; colon and moderately expressed in the bone marrow and stomach. OLFM4 is an antiapoptotic factor that promotes tumor growth. It expressed at high levels in stomach cancer and colon cancer tissues. it promotes proliferation of pancreatic cancer cells by favoring the transition from the S to G2/M phase. In myeloid leukemic cell lines; OLFM4 inhibits cell growth and induces cell differentiation and apoptosis. Through interaction with cell surface lectins and cadherin; OLFM4 facilitates cell adhesion. It may play a role in the inhibition of EIF4EBP1 phosphorylation/deactivation. Induction of OLFM4 in cancer cells was reported to

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com

Catalog Number: PKSH033360



have a novel antiapoptotic action via binding to the potent apoptosis inducer GRIM-19.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com