

Recombinant Human UBE2D1 Protein (GST Tag)

Catalog Number:PKSH033367



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Ubiquitin-conjugating enzyme E2 D1;Stimulator of Fe transport;SFT;UBC4/5 homolog;UbcH5;Ubiquitin carrier protein D1;Ubiquitin-conjugating enzyme E2(17)KB 1;Ubiquitin-conjugating enzyme E2-17 kDa 1;Ubiquitin-protein ligase D1;SFT;UBC5A;UBCH5;UBCH5A
Species	Human
Expression Host	E.coli
Sequence	Met 1-Met 147
Accession	P51668
Calculated Molecular Weight	42.9 kDa
Observed molecular weight	40 kDa
Tag	N-GST

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
Formulation	Supplied as a 0.2 µm filtered solution of 50mM HEPES, 150mM NaCl, 2mM DTT, 10% Glycerol, pH 7.5.
Reconstitution	Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Ubiquitin-conjugating enzyme E2 D1 (UBE2D1) belongs to the ubiquitin-conjugating enzyme family. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes; or E1s; ubiquitin-conjugating enzymes; or E2s; and ubiquitin-protein ligases; or E3s. This enzyme is closely related to a stimulator of iron transport (SFT); and is up-regulated in hereditary hemochromatosis. It also functions in the ubiquitination of the tumor-suppressor protein p53 and the hypoxia-inducible transcription factor HIF1alpha by interacting with the E1 ubiquitin-activating enzyme and the E3 ubiquitin-protein ligases.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017