Recombinant Human UBE2D1 Protein (GST Tag)

Catalog Number: PKSH033367



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Ubiquitin-conjugating enzyme E2 D1;Stimulator of Fe transport;SFT;UBC4/5

homolog;UbcH5;Ubiquitin carrier protein D1;Ubiquitin-conjugating enzyme E2(17)KB 1;Ubiquitin-conjugating enzyme E2-17 kDa 1;Ubiquitin-protein ligase

D1;SFT;UBC5A;UBCH5;UBCH5A

SpeciesHumanExpression HostE.coli

Sequence Met 1-Met 147

Accession P51668
Calculated Molecular Weight 42.9 kDa
Observed molecular weight 40 kDa
Tag N-GST

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Storage Store at $< -20^{\circ}$ C, stable for 6 months. Please minimize freeze-thaw cycles.

Shipping This product is provided as liquid. It is shipped at frozen temperature with blue


ice/gel packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 μm filtered solution of 50mM HEPES, 150mM NaCl, 2mM DTT,

10% Glycerol, pH 7.5.

Reconstitution Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Ubiquitin-conjugating enzyme E2 D1 DBE2D1 belongs to the ubiquitin-conjugating enzyme family. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes; or E1s; ubiquitin-conjugating enzymes; or E2s; and ubiquitin-protein ligases; or E3s. This enzyme is closely related to a stimulator of iron transport (SFT); and is upregulated in hereditary hemochromatosis. It also functions in the ubiquitination of the tumor-suppressor protein p53 and the hypoxia-inducible transcription factor HIF1alpha by interacting with the E1 ubiquitin-activating enzyme and the E3 ubiquitin-protein ligases.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com
Email: techsupport@elabscience.com