

## Recombinant Human GPA33 Protein (His Tag)

Catalog No. PKSH033373

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	Cell Surface A33 Antigen;Glycoprotein A33;GPA33
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Ile22-Val235
<b>Accession</b>	Q99795
<b>Calculated Molecular Weight</b>	24.7 kDa
<b>Observed molecular weight</b>	34-37 kDa
<b>Tag</b>	C-His
<b>Bioactivity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, 1mM EDTA, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Human Glycoprotein A33 (GPA33) is a single-pass type I membrane protein; belongs to the CTX family of cell adhesion

### For Research Use Only

molecular within the immunoglobulin family; can be expressed in normal gastrointestinal epithelium and in 95% of colon cancers. GPA33 consists of one Ig-like C2-type domain and one Ig-like V-type domain. The predicted mature protein includes a single transmembrane domain; a extracellular region and a intracellular tail. Intracellular traffic and recycling to the cell surface appear to play an important role in GPA33 function and to have an influence on its surface density superseding translation regulation. GPA33 has become a promising target of immunologic therapy strategies. GPA33 may also play a important role in cell-cell recognition and signaling.