## Recombinant Human TIGIT Protein (aa 22-141, His Tag)

Catalog Number: PKSH033394



Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

**Synonyms** T-cell immunoreceptor with Ig and ITIM domains;;VSIG9;VSTM3;TIGIT;V-set and

transmembrane domain-containing protein 3;V-set and immunoglobulin domain-

containing protein 9

**Species** Human

HEK293 Cells **Expression Host** Met22-Pro141 **Sequence** Accession Q495A1 Calculated Molecular Weight 14.1 kDa Observed molecular weight 13-19 kDa C-His

#### **Properties**

Tag

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per ug of the protein as determined by the LAL method.

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to **Storage** 

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Please refer to the printed manual for detailed information. Reconstitution

#### Data



> 95 % as determined by reducing SDS-PAGE.

### **Background**

T cell immunoreceptor with Ig and ITIM domains (TIGIT) is a member of the CD28 family within the Ig superfamily of proteins. TIGIT is expressed on NK cells and subsets of activated; memory and regulatory T cells; and particularly on follicular helper T cells within secondary lymphoid organs. It binds to CD155 and Nectin-2 that appear on dendritic cells (DC) and endothelium. Ligation of TIGIT on T cells down-regulates TCR-mediated activation and subsequent proliferation; while NK cell TIGIT ligation blocks NK cell cytotoxicity. Through CD155 and Nectin-2; which also interact with DNAM-1/CD226 and CD96/Tactile; TIGIT is part of an interacting network of Ig superfamily members that may

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

# Recombinant Human TIGIT Protein (aa 22-141, His Tag)

Catalog Number:PKSH033394



augment or oppose each other. In particular; TIGIT binding to CD155 can antagonize the effects of DNAM1.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>