Recombinant Human GFRRA2 Protein (His Tag)

Catalog Number: PKSH033430



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Synonyms	GDNF Family Receptor Alpha-2;GDNF Receptor Alpha-2;GDNFR-Alpha-2; Alpha-2;GDNF Receptor Beta;GDNFR-Beta;Neurturin Receptor Alpha;NRTM Alpha;NTNR-Alpha;RET Ligand 2;TGF-Beta-Related Neurotrophic Factor Receptor 2;GFRA2;GDNFRB;RETL2;TRNR2		
Species	Human		
Expression Host	HEK293 Cells		
Sequence	Ser22-Ser441		
Accession	O00451		
Calculated Molecular Weight	47.8 kDa 80 kDa		
Observed molecular weight			
Tag	C-His		
Properties			
Purity	> 95 % as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method. Generally, lyophilized proteins are stable for up to 12 months when stored at -20 -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliqu of reconstituted samples are stable at < -20°C for 3 months.		
Storage			
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed man		

kDa	MK	R	
120 90	Torigina a	And in case of the local division of the loc	
60		-	
40			
30	-		
20	-		
14	-		

> 95 % as determined by reducing SDS-PAGE.

Background

Members of the glial cell line-derived neurotrophic factor (GDNF) family, including GDNF and Neurturin, play key roles in the control of vertebrate neuronal survivial and differentiation. GDNF is a glycosylated, disulfide-bonded homodimer that is distantly related to the TGF superfamily of growth factors. Three receptors for these factors, GFR α -1, GFR α -2, and GFR α -3 have been identified. The receptors do not contain transmembrane domains and are attached to the cell membrane by glycosyl-phosphoinositol linkage. Both GFR α -1 and GFR α -2 have been shown to mediate the GDNF-

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Recombinant Human GFRRA2 Protein (His Tag)

Catalog Number:PKSH033430



dependent and Neurturin-dependent phosphorylation and activation of the tyrosine kinase Ret. GFR-3 is expressed only during development. GFR α -2 binds Neurturin and mediates activation of RET receptor tyrosine kinase by both Neurturin and GDNF.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com