

# Recombinant Human Fas/CD95/TNFRSF6 Protein (His Tag)



Catalog Number:PKSH033436

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	Tumor necrosis factor receptor superfamily member 6;Apo-1 antigen;Apoptosis-mediating surface antigen FAS;FASLG receptor;APT1;FAS1;TNFRSF6 and FAS;ALPS1A;APO-1;APT1;CD95;FASTM
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Gln26-Asn173
<b>Accession</b>	P25445
<b>Calculated Molecular Weight</b>	17.7 kDa
<b>Observed molecular weight</b>	22-35 kDa
<b>Tag</b>	C-His

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed man
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

FAS is a receptor and contains three TNFR-Cys repeats and one death domain. It has been shown that FAS is involved in the physiological regulation of programmed cell death, and has been implicated in the pathogenesis of various malignancies and diseases of the immune system. FADD (adapter molecule) recruits caspase-8 to the activated receptor, the resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases mediating apoptosis. FAS-mediated apoptosis may play a role in the induction of peripheral tolerance, in the antigen-stimulated suicide of mature T-cells, or both.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017