Recombinant Human IL1R1/CD121a Protein (His Tag)

Catalog Number: PKSH033447



Note: Centrifuge before opening to ensure complete recovery of vial contents.

T	4.0	
Descri	ntian	
DUSCII		

Synonyms CD121A;D2S1473;IL-1 RI;IL-1R-alpha;IL-1R1;IL1R;IL1RA;P80;Interleukin-1

receptor type 1;IL-1R-1;IL-1RT-1;IL-1RT1;CD121 antigen-like family member

A;Interleukin-1 receptor alpha;IL-1R-alpha

Species Human

Expression Host HEK293 Cells **Sequence** Leu18-Thr332

AccessionP14778Calculated Molecular Weight37.0 kDaObserved molecular weight48-66 kDaTagC-His

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Interleukin 1 receptor, type I (IL-1R1) is an interleukin receptor that belongs to the interleukin-1 receptor family. IL-1R1 is an 80 kDa transmembrane protein that is expressed predominantly by T cells, fibroblasts, and endothelial cells. This gene along with IL1R2, IL1RL2, and IL1RL1 form a cytokine receptor gene cluster in a region mapped to chromosome 2q12. IL-1R1 is an important mediator involved in many cytokine induced immune and inflammatory responses. It binds to interleukin-1 associates with the corecptor IL1RAP to form the high affinity interleukin-1 receptor complex which mediates interleukin-1-dependent activation of NF-kappa-B, MAPK and other pathways. The signaling involves the

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

Recombinant Human IL1R1/CD121a Protein (His Tag)

Catalog Number:PKSH033447



recruitment of adapter molecules such as TOLLIP, MYD88, and IRAK1 or IRAK2 via the respective TIR domains of the receptor/coreceptor subunits. It also binds ligands with comparable affinity and binding of antagonist IL1RN prevents association with IL1RAP to form a signaling complex. An IL1 receptor accessory protein that can heterodimerize with the Type I receptor in the presence of IL1 α or IL1 β but not IL1ra, was identified. Recombinant IL1 soluble receptor Type I is a potent antagonist of IL1 action.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>