Recombinant Human IL1R2/CD121b Protein (His Tag)

Catalog Number: PKSH033449



Note: Centrifuge before opening to ensure complete recovery of vial contents.

	_ ·
Descri	ntion

CD121b;CDw121b;IL-1R-2;IL-1RT-2;IL-1RT2;IL1R2c;IL1RB;Interleukin-1 **Synonyms**

receptor type 2;IL-1R-2;CD121 antigen-like family member B;CDw121b;IL-1 type

II receptor;Interleukin-1 receptor beta;IL-1R-beta;Interleukin-1 receptor type I

Species Human

HEK293 Cells **Expression Host** Phe14-Glu343 **Sequence**

Accession P27930 Calculated Molecular Weight 38.6 kDa Observed molecular weight 50 kDa C-His Tag

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per ug of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Please refer to the printed manual for detailed information. Reconstitution

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Interleukin-1 receptor type 2 (IL1R2) belongs to the interleukin-1 receptor family. Two distinct types of IL1 receptors which are able to bind IL1 specifically have been identified; designated as IL1RI (IL1RA) and IL1RII (IL1RB). IL1 receptor type II is a 68 kDa transmembrane protein found on B lymphocytes; neutrophils; monocytes; large granular leukocytes and endothelial cells. IL1R2 is non-signaling receptor forIL1A; IL1B and IL1RN; reduces IL1B activities. IL1R2 serves as a decoy receptor by competetive binding to IL1B and preventing its binding to IL1R1. IL1R2 modulates cellular response through non-signaling association with IL1RAP after binding to IL1B. IL1R2 (membrane and secreted

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

Recombinant Human IL1R2/CD121b Protein (His Tag)

Catalog Number:PKSH033449



forms) preferentially binds IL1B and poorly IL1A and IL1RN. The secreted IL1R2 recruits secreted IL1RAP with high affinity; this complex formation may be the dominant mechanism for neutralization of IL1B by secreted/soluble receptors.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>