

# Recombinant Mouse FCRL1 Protein (aa 17-219, His Tag)

Catalog Number:PKSM041196



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

|                                    |  |
|------------------------------------|--|
| <b>Synonyms</b>                    | Fc receptor-like protein 1;FcR-like protein 1;FcRL1;BXMAS1-like protein 1;mBXMH1;Fc receptor homolog 1;FcRH1;moFcRH1;IFGP family protein 1;mIFGP1;CD307a;Bxmas1;Bxmh1;Bxmh1b;Fcrh1;Fcrh1l;Fcrh1s;Ifgp1 Protein |
| <b>Species</b>                     | Mouse  |
| <b>Expression Host</b>             | HEK293 Cells   |
| <b>Sequence</b>                    | Ala17-Ser219   |
| <b>Accession</b>                   | BAC30017.1   |
| <b>Calculated Molecular Weight</b> | 22.5 kDa   |
| <b>Observed molecular weight</b>   | 40-60 kDa  |
| <b>Tag</b>                         | C-His  |

## Properties

|                       |   |
|-----------------------|---|
| <b>Purity</b>         | > 95 % as determined by reducing SDS-PAGE.  |
| <b>Endotoxin</b>      | < 1.0 EU per µg of the protein as determined by the LAL method.   |
| <b>Storage</b>        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.             |
| <b>Shipping</b>       | This product is provided as lyophilized powder which is shipped with ice packs.   |
| <b>Formulation</b>    | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.<br>Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual. |
| <b>Reconstitution</b> | Please refer to the printed manual for detailed information.  |

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Mouse Fc receptor-like protein 1 (FCRL1) is a single-pass type I membrane protein. It is expressed in B-cells at the various stages of differentiation. Mouse FCRL1 consists of a 203 amino acid (aa) extracellular domain (ECD) with two Ig-like domains, a 21 aa transmembrane segment, and a 103 aa cytoplasmic domain with two immunotyrosine activation motifs (ITAMs). FCRL1 is likely not a receptor for immunoglobulin. It may function as an activating coreceptor in B-cells and B-cells differentiation.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017