

## Recombinant Mouse Semaphorin-4D/SEMA4D Protein (Fc Tag)

**Catalog No.** PKSM041209

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	SEMA4D;Semaphorin-4D;M-Sema G;Semaphorin-C-like 2;Semaphorin-J;Sema J;CD100;Semacl2;Semaj;coll-4;Semacl2;Semaj
<b>Species</b>	Mouse
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Phe24-Met711
<b>Accession</b>	O09126
<b>Calculated Molecular Weight</b>	103.4 kDa
<b>Observed molecular weight</b>	110-130 kDa
<b>Tag</b>	C-Fc
<b>Bioactivity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 10% Trehalose, 100mM NaCl, 0.05% Tween 80, pH 7.5. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

SEMA4D is a member of the semaphorin family, contains one Ig-like C2-type domain, one PSI domain and one Sema domain. SEMA4D is strongly expressed in lymphoid tissues, especially in the thymus, as well as in the nervous tissues. However, SEMA4D is expressed at lower levels in testes, brain, kidney, small intestine, prostate, heart, placenta, lung and pancreas, but not in colon and liver. SEMA4D is a cell surface receptor for PLXN1B and PLXNB2 that plays an important role in cell-cell signaling. SEMA4D is involved in a number of fundamental biological processes such as promoting reorganization of the actin cytoskeleton, the migration of cerebellar granule cells and of endothelial cells and signaling via SRC and PTK2B/PYK2, which then mediates activation of phosphatidylinositol 3-kinase and of the AKT1 signaling cascade. Not only these, it plays a role in axonal growth cone guidance in the developing central nervous system. Semaphorin-4D / SEMA4D may play a functional role in the immune system, as well as in the nervous system. It could induce B-cells to aggregate and improves their viability (in vitro). SEMA4D is involved in regulating dendrite and axon branching and morphogenesis and promoting interaction with PLXNB1 mediates activation of RHOA.