## **Recombinant Mouse TNFR1/TNFRSF1A Protein**

Catalog No. PKSM041215

*Note:* Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Tumor necrosis factor receptor superfamily member 1A;Tumor necrosis factor receptor 1;Tumor necrosis factor receptor type I;Tnfr-1;Tnfr1;Tnfrsf1a;
Species	Mouse
Expression Host	E.coli
Sequence	Ile22-Ala212
Accession	P25118
Calculated Molecular Weight	21.2 kDa
Observed molecular weight	20-27 kDa
Tag	None
Bioactivity	Not validated for activity
Properties	
Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.5. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	



> 90 % as determined by reducing SDS-PAGE.

## Background

Tumor necrosis factor receptor superfamily member 1A (Tnfrsf1a) is a member of the tumor necrosis factor receptor

## For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u>

## **Elabscience**®

superfamily. Tnfrsf1a is one of the major receptors for the tumor necrosis factor-alpha. It can activate the transcription factor NF- $\kappa$ B, mediate apoptosis, and function as a regulator of inflammation. Antiapoptotic protein BCL2-associated athanogene 4 (BAG4/SODD) and adaptor proteins TRADD and TRAF2 have been shown to interact with this receptor, and thus play regulatory roles in the signal transduction mediated by the receptor. Germline mutations of the extracellular domains of this receptor were found to be associated with the human genetic disorder called tumor necrosis factor associated periodic syndrome (TRAPS) or periodic fever syndrome

For Research Use Only