

# Recombinant Mouse CD226/DNAM-1 Protein (His Tag)

Catalog Number:PKSM041233



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	CD226 antigen;platelet and T cell activation antigen 1;CD226 molecule;DNAM1 adhesion glycoprotein;DNAM-1;DNAX accessory molecule-1;DNAX accessory molecule 1;PTA1;T lineage-specific activation antigen 1 antigen;CD226;PTA1;TLISA1;BC051526;DNAM-1;DNAM1;Pta1;TLISA1
<b>Species</b>	Mouse
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Glu19-Pro254
<b>Accession</b>	Q8K4F0
<b>Calculated Molecular Weight</b>	27.6 kDa
<b>Observed molecular weight</b>	35-50 kDa
<b>Tag</b>	C-His

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Mouse DNAX accessory molecule-1(DNAM-1) is a type I transmembrane glycoprotein that belongs to the immunoglobulin superfamily. As an activating receptor, it interacts with the ligands CD155 and CD112, and activates natural killer (NK) cells via its immunoreceptor tyrosine-based activatory motif (ITAM). Mature mouse DNAM-1 has extracellular domain (ECD) that contains two Ig-like C2-set domains, and possesses a cytoplasmic region that contains motifs for binding PDZ domains. DNAM-1 is expressed on several lymphoid and myeloid cell types and interacts with

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Mouse CD226/DNAM-1 Protein (His Tag)

Catalog Number:PKSM041233



CD155/PVR and Nectin-2/CD112. Ligation of DNAM-1 promotes the activation of NK cells, CD8+ T cells, and mast cells, induces dendritic cell maturation, initiates megakaryocyte and activated platelet adhesion to vascular endothelial cells, and stimulates monocyte extravasation? Conversely, it inhibits the formation of osteoclasts. Platelet-endothelium interactions that are mediated by DNAM-1 enable the metastasis of tumor cells to the lung.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)