

Recombinant Mouse BMPRIA/ALK-3 Protein (Fc & His Tag)

Catalog No. PKSM041249

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|------------------------------------|--|
| Synonyms | ALK-3;Bone morphogenetic protein receptor type-1A;BMP type-1A receptor;BMPR-1A;Activin receptor-like kinase 3;BMP-2/BMP-4 receptor;Serine/threonine-protein kinase receptor R5;SKR5;CD292;Acvrlk3;Bmpr;BMPR-IA |
| Species | Mouse |
| Expression Host | HEK293 Cells |
| Sequence | Gln24-Arg152 |
| Accession | P36895 |
| Calculated Molecular Weight | 42.2 kDa |
| Observed molecular weight | 55-60&120 kDa |
| Tag | C-Fc-His |
| Bioactivity | Not validated for activity |

Properties

| | |
|-----------------------|--|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 95 % as determined by reducing SDS-PAGE.

For Research Use Only

Background

ALK-3 is a type I receptor for bone morphogenetic proteins (BMPs) which belong to the protein kinase superfamily, TKL Ser/Thr protein kinase family and TGFB receptor subfamily. The BMP receptors consists of the type I receptors BMPR1A and BMPR1B and the type I I receptor BMPR2. Seven known type I serine/threonine kinases and five mammalian type II serine/threonine kinase receptors function in TGF-beta superfamily signal transduction. The downstream molecules of the type I BMP receptors include the Smad (Smad1, 5 and 8) proteins that are phosphorylated in a ligand-dependent manner, and relay the BMP signal from the receptors to target genes in the nucleus. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then bind and activate SMAD transcriptional regulators. ALK-3 contains a GS domain and a protein kinase domain. ALK-3 is widely expressed. Defects in BMPR1A gene are a cause of a significant proportion of cases of Juvenile polyposis syndrome (JPS).

For Research Use Only