



A Reliable Research Partner in Life Science and Medicine

## Purified Anti-Mouse CD279/PD-1 Antibody[29F.1A12]

Catalog No.E-AB-F1131AReactivityMouseStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsFCM

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### **Antigen Information**

Alternate Names PD-1, Programmed Death-1

**Uniprot ID** Q02242

**Background** CD279, also known as programmed death-1 (PD-1), is a 50-55 kD glycoprotein belonging to the

CD28 family of the Ig superfamily. PD-1 is expressed on activated splenic T and B cells and thymocytes. It is induced on activated myeloid cells as well. PD-1 is involved in lymphocyte clonal selection and peripheral tolerance through binding its ligands, B7-H1 (PD-L1) and B7-DC (PD-L2). It has been reported that PD-1 and PD-L1 interactions are critical to positive selection and play a role in shaping the T cell repertoire. PD-L1 negative costimulation is essential for

prolonged survival of intratesticular islet allografts.

#### **Product Details**

 $\begin{tabular}{lll} Form & Liquid \\ Concentration & 0.5 mg/mL \\ Size & 25 \mu g/100 \mu g \\ Clone No. & 29F.1A12 \\ Host & Rat \\ \end{tabular}$ 

IsotypeRat IgG2a, κReactivityMouseApplicationFCM

Isotype ControlPurified Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833A]Storage BufferPhosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Shipping
Biological ice pack at 4 °C
Stability & Storage
Keep as concentrated solution.
Store at 2~8°C .Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: <a href="https://www.elabscience.com">www.elabscience.com</a>
 Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>

#### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

# **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 0.25 \,\mu g$  per  $10^6$  cells in 100  $\mu L$  volume or 100  $\mu L$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

### **Related Information**

- 1. Sample Preparation for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5594.html">https://www.elabscience.com/List-detail-5594.html</a>
- 2. Staining Cell Surface Targets for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5568.html">https://www.elabscience.com/List-detail-5568.html</a>
- 3. Flow Cytometry Troubleshooting Tips <a href="https://www.elabscience.com/List-detail-5593.html">https://www.elabscience.com/List-detail-5593.html</a>
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/Listdetail-459742.html

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com

Fax: 1-832-243-6017