



A Reliable Research Partner in Life Science and Medicine

APC Anti-Human IL-4 Antibody[MP4-25D2]

Catalog No.E-AB-F1203EReactivityHumanStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsICFCM

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names Interleukin-4,IL-4,B-cell IgG differentiation factor,B-cell growth factor 1,BSF-1,IGG1 induction

factor

Uniprot ID P05112

Background IL-4 is a pleiotropic cytokine that is produced by activated T cells, mast cells, and basophils. IL-4

elicits many different biological responses but has two dominant functions. The first is regulating differentiation of naïve CD4+ T cell to the Th2 type. Th2 cells produce IL-4, IL-5, IL-10, and IL-13, which tend to favor a humoral immune response while suppressing a cell-mediated

immune response controlled by Th1 cells. The second is regulating IgE and IgG1 production by B

cells.

Product Details

Form Liquid

Size 20Tests/100Tests/100Tests×2

Clone No. MP4-25D2

Host Rat

 $\begin{tabular}{lll} \textbf{Isotype} & Rat IgG1, \kappa \\ \textbf{Reactivity} & Human \\ \textbf{Application} & ICFCM \\ \end{tabular}$

Isotype Control APC Rat IgG1, κ Isotype Control[HRPN] [Product E-AB-F09822E]

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping Biological ice pack at 4 °C **Stability & Storage** Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

Elabscience Bionovation Inc.

Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

Fluorophore

Conjugation: APC

APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Intracellular Antigens for Flow Cytometry https://www.elabscience.com/List-detail-5570.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/List-detail-459742.html

For Research Use Only

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086