



A Reliable Research Partner in Life Science and Medicine

### APC Anti-Mouse IL-4 Antibody[11B11]

Catalog No. E-AB-F1204E Reactivity Mouse Storage Store at 2~8°C, Avoid freeze / thaw cycles **Applications ICFCM** 

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### **Antigen Information**

**Alternate Names** Interleukin-4,IL-4,B-cell IgG differentiation factor,B-cell growth factor 1,BSF-1,IGG1 induction

**Uniprot ID** P07750

**Background** IL-4 is a pleiotropic cytokine produced by activated T cells, mast cells, and basophils. IL-4 is a

potent lymphoid cell growth factor which stimulates the growth and activation of certain B cells

and T cells. IL-4 is important for regulation of T helper subset development.

#### **Product Details**

Form Liquid

50Tests/100Tests/100Tests×2 Size

Clone No. 11B11 Host Rat **Isotype** Rat IgG1, κ Reactivity Mouse **Application ICFCM** 

**Isotype Control** APC Rat IgG1, κ Isotype Control[HRPN] [Product E-AB-F09822E]

**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping** Biological ice pack at 4 °C Stability & Storage Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com

Email: techsupport@elabscience.com

A Reliable Research Partner in Life Science and Medicine

## **Fluorophore**


Conjugation: APC

APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).

# **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### **Product data**



HEK293T cells transiently transfected with pcDNA3.1 plasmid encoding Mouse IL-4 gene are stained with APC Anti-Mouse IL-4 Antibody (filled gray histogram) or APC Rat IgG1,  $\kappa$  Isotype Control (empty black histogram).

#### **Related Information**

- 1. Sample Preparation for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5594.html">https://www.elabscience.com/List-detail-5594.html</a>
- 2. Staining Intracellular Antigens for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5570.html">https://www.elabscience.com/List-detail-5570.html</a>
- 3. Flow Cytometry Troubleshooting Tips <a href="https://www.elabscience.com/List-detail-5593.html">https://www.elabscience.com/List-detail-5593.html</a>
- 4. How to select the appropriate detection channel through the spectrogram? <a href="https://www.elabscience.com/List-detail-459742.html">https://www.elabscience.com/List-detail-459742.html</a>

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086

 Web: <a href="www.elabscience.com">www.elabscience.com</a>
 Email: <a href="techsupport@elabscience.com">techsupport@elabscience.com</a>