

## PE Anti-Human/Mouse IL-5 Antibody[TRFK5]

<b>Catalog No.</b>	E-AB-F1205D	<b>Reactivity</b>	Human,Mouse
<b>Storage</b>	Store at 2~8°C, Avoid freeze / thaw cycles	<b>Applications</b>	ICFCM

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Antigen Information

<b>Alternate Names</b>	Interleukin-5,IL-5,B-cell differentiation factor I,T-cell replacing factor,TRF
<b>Uniprot ID</b>	P05113,P04401
<b>Background</b>	IL-5 is a homodimeric, disulphide-linked protein produced by T-cells. Monomeric human IL-5 is a 126 amino acid protein with a reported molecular weight of 26 kD for the homodimeric protein. Mouse and human IL-5 are approximately 70% identical. IL-5 has been shown to promote the growth of immature hematopoietic BFU-E progenitors, stimulate the activation and differentiation of eosinophils, and promote the generation of cytotoxic lymphocytes. Mouse IL-5 induces the differentiation and proliferation of pre-activated B-cells and stimulates the production and secretion of IgM and IgA by B-cells stimulated with bacterial endotoxin. The TRFK5 antibody can neutralize the bioactivity of natural or recombinant IL-5.

### Product Details

<b>Form</b>	Liquid
<b>Size</b>	20Tests/100Tests/100Tests×2
<b>Clone No.</b>	TRFK5
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG1, κ
<b>Reactivity</b>	Human,Mouse
<b>Application</b>	ICFCM
<b>Isotype Control</b>	<a href="#">PE Rat IgG1, κ Isotype Control[HRPN] [Product E-AB-F09822D]</a>
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
<b>Shipping</b>	Biological ice pack at 4 °C
<b>Stability &amp; Storage</b>	Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light.Do not freeze. This product is guaranteed up to one year from purchase.

### For Research Use Only

## Fluorophore


### Conjugation: PE

PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green (561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42 nm bandpass filter).

## Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## Product data



CHO cells transiently transfected with pcDNA3.1 plasmid encoding Mouse IL-5 gene are stained with PE Anti-Human/Mouse IL-5 Antibody[TRFK5] (filled gray histogram) or PE Rat IgG1,  $\kappa$  Isotype Control (empty black histogram).

## Related Information

1. Sample Preparation for Flow Cytometry <https://www.elabscience.com/List-detail-5594.html>
2. Staining Intracellular Antigens for Flow Cytometry <https://www.elabscience.com/List-detail-5570.html>
3. Flow Cytometry Troubleshooting Tips <https://www.elabscience.com/List-detail-5593.html>
4. How to select the appropriate detection channel through the spectrogram? <https://www.elabscience.com/List-detail-459742.html>