



A Reliable Research Partner in Life Science and Medicine

# AF/LE Purified Anti-Mouse CD25 Antibody[PC-61.5.3]

Catalog No.E-AB-F11020ReactivityMouseStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsFCM,Block

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### **Antigen Information**

Alternate Names Interleukin-2 receptor subunit alpha,IL-2RA,IL-2 receptor subunit alpha,IL-2-RA,IL-2R subunit

alpha,IL2-RA,TAC antigen,p55,p55

Uniprot ID P01590

**Background** CD25 is a 55 kD glycoprotein, also known as the low affinity IL-2Rα, Ly-43, p55, or Tac. It is

expressed on activated T and B cells, thymocyte subset, pre-B cells, and T regulatory cells. In association with CD122 (IL-2R $\beta$ ) and CD132(common  $\gamma$  chain), CD25 forms the high affinity

signaling IL-2 receptor.

#### **Product Details**

 Form
 Liquid

 Concentration
 0.5 mg/mL

 Size
 50μg/500μg/1mg

 Clone No.
 PC-61.5.3

Clone No. PC-6
Host Rat

 $\begin{tabular}{lll} \textbf{Isotype} & Rat IgG1, \kappa \\ \textbf{Reactivity} & Mouse \\ \textbf{Application} & FCM,Block \\ \end{tabular}$ 

**Isotype Control** AF/LE Purified Rat IgG1, κ Isotype Control[HRPN] [Product E-AB-F098230]

Storage Buffer 0.2 µm filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or

stabilizers. Endotoxin level is < 2 EU/mg as Determined by LAL gel clotting assay.

**Shipping** Biological ice pack at 4 °C **Stability & Storage** Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com





A Reliable Research Partner in Life Science and Medicine

## **Fluorophore**

**Conjugation:** None (Purified antibody-Azide Free/Low endotoxin)

#### Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 2.0 \,\mu g$  per  $10^6$  cells in  $100 \,\mu L$  volume or  $100 \,\mu L$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

#### **Related Information**

- 1. Sample Preparation for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5594.html">https://www.elabscience.com/List-detail-5594.html</a>
- 2. Staining Cell Surface Targets for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5568.html">https://www.elabscience.com/List-detail-5568.html</a>
- 3. Flow Cytometry Troubleshooting Tips <a href="https://www.elabscience.com/List-detail-5593.html">https://www.elabscience.com/List-detail-5593.html</a>
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/Listdetail-459742.html

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com

Fax: 1-832-243-6017