



A Reliable Research Partner in Life Science and Medicine

AF/LE Purified Anti-Mouse CD3ε Antibody[145-2C11]

Catalog No. E-AB-F11030 Reactivity Mouse Storage Store at 2~8°C, Avoid freeze / thaw cycles **Applications** FCM, Activ

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names T-cell surface glycoprotein CD3 epsilon chain, CD3E, T-cell surface antigen T3/Leu-4 epsilon

chain,CD3e,CD3E,T3E

Uniprot ID P22646

Background CD3 is a 20 kD transmembrane protein, also known as CD3 or T3. It is a member of the Ig

> superfamily and primarily expressed on T cells, NK-T cells, and at different levels on thymocytes during T cell differentiation. CD3ε forms a TCR complex by associating with the CD3δ, γ and ζ chains, as well as the TCR α/β or γ/δ chains. CD3 plays a critical role in TCR signal transduction,

T cell activation, and antigen recognition by binding the peptide/MHC antigen complex.

Product Details

Form Liquid Concentration 0.5 mg/mLSize $50 \mu g / 500 \mu g / 1 mg$ Clone No. 145-2C11

Host Armenian Hamster Armenian Hamster IgG **Isotype**

Reactivity Mouse **Application** FCM.Activ

Isotype Control AF/LE Purified Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F098530]

Storage Buffer 0.2 µm filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or

stabilizers. Endotoxin level is < 2 EU/mg as Determined by LAL gel clotting assay.

Shipping Biological ice pack at 4 °C Stability & Storage Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com

Email: techsupport@elabscience.com





A Reliable Research Partner in Life Science and Medicine

Fluorophore

Conjugation: None (Purified antibody-Azide Free/Low endotoxin)

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 2.0 \,\mu g$ per 10^6 cells in $100 \,\mu L$ volume or $100 \,\mu L$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/Listdetail-459742.html

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com

Fax: 1-832-243-6017