



A Reliable Research Partner in Life Science and Medicine

AF/LE Purified Anti-Mouse Ly6C Antibody[Monts 1]

Catalog No.E-AB-F11210ReactivityMouseStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsFCM

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names Lymphocyte antigen 6 complex, locus C,Ly 6C,Ly6c1,Ly6c2

Uniprot ID P0CW03

Background Most hematopoietic cells express one or more members of Ly-6 family. The expression of Ly-6

varies with development stage and activation. Ly-6C is a 14-17 kD GPI-linked surface protein expressed on mouse monocyte/macrophage cells, endothelial cells, neutrophils, and some T cell

subsets. Ly-6C is reported to be an indicator of memory CD8+ T cells.

Product Details

 $\begin{array}{lll} \textbf{Form} & & \text{Liquid} \\ \textbf{Concentration} & & 0.5 \text{ mg/mL} \\ \textbf{Size} & & 50 \mu \text{g/}500 \mu \text{g/}1 \text{mg} \\ \end{array}$

Clone No. Monts 1
Host Rat

 $\begin{tabular}{lll} \textbf{Isotype} & Rat IgG2a, \kappa \\ \textbf{Reactivity} & Mouse \\ \textbf{Application} & FCM \\ \end{tabular}$

Isotype Control AF/LE Purified Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F098330]

Storage Buffer 0.2 µm filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or

stabilizers. Endotoxin level is < 2 EU/mg as Determined by LAL gel clotting assay.

Shipping Biological ice pack at 4 °C **Stability & Storage** Keep as concentrated solution.

Store at $2\sim8^{\circ}\text{C}$ and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com



Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

Fluorophore

Conjugation: None (Purified antibody-Azide Free/Low endotoxin)

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \,\mu g$ per 10^6 cells in $100 \,\mu L$ volume or $100 \,\mu L$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/Listdetail-459742.html

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com