



A Reliable Research Partner in Life Science and Medicine

AF/LE Purified Anti-Human CD44 Antibody[Hermes-1]

Catalog No. E-AB-F12150 Reactivity Human Storage Store at 2~8°C, Avoid freeze / thaw cycles **Applications** Block.FCM

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names LHR,PGP-I,CDw44,Epican,HUTCH-I

Uniprot ID P16070

Background CD44 is a 80-95 kD glycoprotein also known as Hermes, Pgp1, H-CAM, or HUTCH. It is

expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. As B and T cells become activated or progress to the memory stage, CD44 expression increases from a low or mid level of intensity to high expression levels. Thus, CD44 has been reported to be a valuable marker for memory cell subsets. CD44 is an adhesion molecule involved in leukocyte attachment to and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of

inflammation, and leukocyte aggregation.

Product Details

Form Liquid Concentration 0.5 mg/mL50μg/500μg/1mg Size

Clone No. Hermes-1 Host Rat

Rat IgG2a, κ **Isotype** Reactivity Human **Application** Block,FCM

Isotype Control AF/LE Purified Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F098330]

0.2 µm filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or **Storage Buffer**

stabilizers. Endotoxin level is < 2 EU/mg as Determined by LAL gel clotting assay.

Shipping Biological ice pack at 4 °C

Stability & Storage Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com

Email: techsupport@elabscience.com

Fax: 1-832-243-6017





A Reliable Research Partner in Life Science and Medicine

Fluorophore

Conjugation: None (Purified antibody-Azide Free/Low endotoxin)

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 2.0 \,\mu g$ per 10^6 cells in $100 \,\mu L$ volume or $100 \,\mu L$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/Listdetail-459742.html

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com

Fax: 1-832-243-6017