

Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

# AF/LE Purified Anti-Mouse H-2 Antibody[M1/42]

Catalog No.E-AB-F12160ReactivityMouseStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsBlock,FCM

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Antigen Information**

Alternate Names Mouse major histocompatibility complex (MHC) H-2, MHC I

**Background** The M1/42 antibody reacts with the H-2 MHC class I alloantigens expressed on nucleated cells

from mice of the a, b, d, j, k, s, and u haplotypes (Stallcup, KC et al, 1981). MHC class I is

involved in antigen presentation to T cells expressing CD3/TCR and CD8 proteins.

#### **Product Details**

 Form
 Liquid

 Concentration
 0.5 mg/mL

 Size
 50µg/500µg/1mg

Clone No. M1/42 Host Rat

 $\begin{tabular}{lll} \textbf{Isotype} & Rat IgG2a, \kappa \\ \textbf{Reactivity} & Mouse \\ \textbf{Application} & Block,FCM \\ \end{tabular}$ 

Isotype Control AF/LE Purified Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F098330]

Storage Buffer 0.2 µm filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or

stabilizers. Endotoxin level is < 2 EU/mg as Determined by LAL gel clotting assay.

**Shipping** Biological ice pack at 4 °C **Stability & Storage** Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086

 Web: <a href="www.elabscience.com">www.elabscience.com</a>
 Email: <a href="techsupport@elabscience.com">techsupport@elabscience.com</a>



Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

# **Fluorophore**

**Conjugation:** None (Purified antibody-Azide Free/Low endotoxin)

# Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 0.25 \,\mu g$  per  $10^6$  cells in 100  $\mu L$  volume or 100  $\mu L$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

### **Related Information**

- 1. Sample Preparation for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5594.html">https://www.elabscience.com/List-detail-5594.html</a>
- 2. Staining Cell Surface Targets for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5568.html">https://www.elabscience.com/List-detail-5568.html</a>
- 3. Flow Cytometry Troubleshooting Tips <a href="https://www.elabscience.com/List-detail-5593.html">https://www.elabscience.com/List-detail-5593.html</a>
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/Listdetail-459742.html

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086