# PE/Cyanine5 Anti-Mouse CD22 Antibody[Cy34.1]

| | | | |
|---|---|---|---|
| **Catalog No.** | E-AB-F1021UG | **Reactivity** | Mouse |
| **Storage** | Store at 2~8℃, Avoid freeze / thaw cycles | **Applications** | FCM |

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Antigen Information

| | |
|---|---|
| **Alternate Names** | B-cell receptor CD22,Cd22,B-lymphocyte cell adhesion molecule,BL-CAM,Sialic acid-binding Ig-like lectin 2,Siglec-2,T-cell surface antigen Leu-14,CD22,Lyb-8, Siglec2 |
| **Uniprot ID** | P35329 |
| **Background** | The Cy34.1 monoclonal antibody specifically binds to the B-lymphocyte differentiation antigen CD22 on strains having the Lyb-8.2 alloantigen (e.g., A, BALB/c, CBA, C3H/He, C57BL, C57L, C58, SJL, SWR, but not AKR, DBA/1, DBA/2, NZB, PL). CD22 is expressed at high levels on mature peripheral B lymphocytes (follicular and marginal zone), B-1 cells (CD5+ B cells), and plasma cells. It is a member of the Ig gene superfamily and associates with the B-cell antigen receptor. Its sialic acid- binding immunoglobulin-like lectin (siglec) extracellular region mediates B-cell adhesion to ligands on endothelial cells in the bone marrow. Its intracellular domain is phosphorylated after cross-linking of antigen receptor or MHC class II antigen. It is involved in negative regulation of B-cell activation and protection from autoimmunity. B-cell proliferative responses to LPS or anti-mouse Ig μ chain are augmented in the presence of Cy34.1 mAb. |

## Product Details

| | |
|---|---|
| **Form** | Liquid |
| **Concentration** | 0.2 mg/mL |
| **Size** | 25μg/100μg |
| **Clone No.** | Cy34.1 |
| **Host** | Mouse |
| **Isotype** | Mouse IgG1, κ |
| **Reactivity** | Mouse |
| **Application** | FCM |
| **Isotype Control** | [Product E-AB-F09793G] |
| **Storage Buffer** | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant. |
| **Shipping** | Biological ice pack at 4 ℃ |
| **Stability & Storage** | Keep as concentrated solution.<br>Store at 2~8℃ and protected from prolonged exposure to light.Do not freeze.<br>This product is guaranteed up to one year from purchase. |

## Fluorophore

**Conjugation:** PE/Cyanine5

PE/Cyanine5 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 670 nm (e.g., a 690/50 nm bandpass filter).

## Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μg/$10^6$ cells in 100 μL volume].

## Related Information

1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/List-detail-459742.html


Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017
Web: www.elabscience.com Email: techsupport@elabscience.com