



A Reliable Research Partner in Life Science and Medicine

Elab Fluor® 647 Anti-Mouse CD106 Antibody[M/K-2.7]

Catalog No.E-AB-F1091UMReactivityMouseStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsFCM

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names Vascular cell adhesion protein 1,Vcam1,V-CAM 1,VCAM-1,CD106

Uniprot ID P2953

Background CD106 is a 110 kD glycosylphosphatidylinositol (GPI)-linked transmembrane protein, also known

as VCAM-1 and INCAM-110. It is constitutively expressed on bone marrow stromal cells, myeloid progenitors, splenic dendritic cells, activated endothelial cells, as well as some lymphocytes. CD106 expression can be upregulated on endothelial cells by inflammatory cytokines. CD106 is involved in adhesion and acts as a counter-receptor for VLA-4 ($\alpha 4/\beta 1$

integrin) and LPAM-1 ($\alpha 4/\beta 7$ integrin).

Product Details

FormLiquidConcentration0.5 mg/mLSize $25 \mu g/100 \mu g$ Clone No.M/K-2.7HostRatIsotypeRat IgG1, κPenetivityMoves

Reactivity Mouse
Application FCM

Isotype Control Elab Fluor[®] 647 Rat IgG1, κ Isotype Control[HRPN] [Product E-AB-F09823M]

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Shipping Biological ice pack at 4 °C **Stability & Storage** Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

A Reliable Research Partner in Life Science and Medicine

Fluorophore


Conjugation: Elab Fluor® 647

Elab Fluor $^{\odot}$ 647 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 670 nm (e.g., a 660/20 nm bandpass filter).

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is $0.1-1~\mu g/10^6$ cells in $100~\mu L$ volume].

Product data



C57BL/6 murine bone marrow cells are stained with Elab Fluor[®] 647 Anti-Mouse CD106 Antibody (filled gray histogram). Unstained bone marrow cells (empty black histogram) are used as control.

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/List-detail-459742.html

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086

 Web: www.elabscience.com
 Email: techsupport@elabscience.com