

Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

# PE Anti-Mouse CD18 Antibody[M18/2]

Catalog No.E-AB-F1113UDReactivityMouseStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsFCM

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Antigen Information**

Alternate Names Integrin beta-2,Itgb2,Cell surface adhesion glycoproteins LFA-1/CR3/p150+95 subunit

beta, Complement receptor C3 subunit beta, CD18

**Uniprot ID** P11835

**Background** CD18 is a 95 kD protein, also known as integrin β2 subunit. It is expressed on all leukocytes.

CD18, in association with integrin  $\alpha$  chain CD11a, CD11b, and CD11c forms LFA-1, Mac-1, and  $\alpha$ X $\beta$ 2, respectively, and plays an important role in leukocytes adhesion. The CD18 integrin complexes bind ICAM-1 (CD54), ICAM-2 (CD102), ICAM-3 (CD50), iC3b, and fibrinogen.

#### **Product Details**

 $\begin{tabular}{lll} Form & Liquid \\ Concentration & 0.2 mg/mL \\ Size & 25 \mu g/100 \mu g \\ Clone No. & M18/2 \\ Host & Rat \\ \end{tabular}$ 

 $\begin{array}{lll} \textbf{Isotype} & \text{Rat IgG2a, } \kappa \\ \textbf{Reactivity} & \text{Mouse} \\ \textbf{Application} & \text{FCM} \\ \end{array}$ 

**Isotype Control** PE Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833D]

**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping** Biological ice pack at 4 °C **Stability & Storage** Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086

#### **Elabscience Bionovation Inc.**

Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

## **Fluorophore**

Conjugation: PE

PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green (561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42 nm bandpass filter).

## **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is  $0.1-1~\mu g/10^6$  cells in  $100~\mu L$  volume].

#### **Related Information**

- 1. Sample Preparation for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5594.html">https://www.elabscience.com/List-detail-5594.html</a>
- 2. Staining Cell Surface Targets for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5568.html">https://www.elabscience.com/List-detail-5568.html</a>
- 3. Flow Cytometry Troubleshooting Tips <a href="https://www.elabscience.com/List-detail-5593.html">https://www.elabscience.com/List-detail-5593.html</a>
- 4. How to select the appropriate detection channel through the spectrogram? <a href="https://www.elabscience.com/List-detail-459742.html">https://www.elabscience.com/List-detail-459742.html</a>

For Research Use Only

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>