

Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

# PerCP/Cyanine5.5 Anti-Mouse Ly6C Antibody[Monts 1]

Catalog No.E-AB-F1121UJReactivityMouseStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsFCM

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### **Antigen Information**

Alternate Names Lymphocyte antigen 6 complex, locus C,Ly 6C,Ly6c1,Ly6c2

Uniprot ID P0CW03

**Background** Most hematopoietic cells express one or more members of Ly-6 family. The expression of Ly-6

varies with development stage and activation. Ly-6C is a 14-17 kD GPI-linked surface protein expressed on mouse monocyte/macrophage cells, endothelial cells, neutrophils, and some T cell

subsets. Ly-6C is reported to be an indicator of memory CD8+ T cells.

#### **Product Details**

 $\begin{tabular}{lll} Form & Liquid \\ Concentration & 0.2 mg/mL \\ Size & 25 \mu g/100 \mu g \\ Clone No. & Monts 1 \\ Host & Rat \\ \end{tabular}$ 

 $\begin{array}{lll} \textbf{Isotype} & \text{Rat IgG2a, } \kappa \\ \textbf{Reactivity} & \text{Mouse} \\ \textbf{Application} & \text{FCM} \\ \end{array}$ 

Isotype Control PerCP/Cyanine 5.5 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833J]

**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping** Biological ice pack at 4 °C **Stability & Storage** Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

A Reliable Research Partner in Life Science and Medicine

## **Fluorophore**


Conjugation: PerCP/Cyanine5.5

PerCP/Cyanine5.5 is designed to be excited by the blue laser (488 nm) and detected using an optical filter centered near 675 nm (e.g., a 690/50 nm bandpass filter).

### **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is  $0.1-1~\mu g/10^6$  cells in  $100~\mu L$  volume].

#### **Product data**



C57BL/6 murine bone marrow cells are stained with PerCP/Cyanine5.5 Anti-Mouse Ly6C Antibody (filled gray histogram). Unstained bone marrow cells (empty black histogram) are used as control.

#### **Related Information**

- 1. Sample Preparation for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5594.html">https://www.elabscience.com/List-detail-5594.html</a>
- 2. Staining Cell Surface Targets for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5568.html">https://www.elabscience.com/List-detail-5568.html</a>
- 3. Flow Cytometry Troubleshooting Tips <a href="https://www.elabscience.com/List-detail-5593.html">https://www.elabscience.com/List-detail-5593.html</a>
- 4. How to select the appropriate detection channel through the spectrogram? <a href="https://www.elabscience.com/List-detail-459742.html">https://www.elabscience.com/List-detail-459742.html</a>

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086

 Web: <a href="www.elabscience.com">www.elabscience.com</a>
 Email: <a href="techsupport@elabscience.com">techsupport@elabscience.com</a>